Roll No.

Total No. of Pages: 02

Total No. of Questions: 19

PIT M.Sc (Chemistry) (Sem.-1) PHYSICAL CHEMISTRY-I (THERMODYNAMICS AND ELECTROCHEMISTRY)

Subject Code: CHL-403 Paper ID: [51142]

Time: 3 Hrs. Max. Marks: 70

INSTRUCTIONS TO CANDIDATES:

- SECTION-A is COMPULSORY consisting of TEN questions carrying TWO marks each.
- 2. SECTION-B contains SIX questions carrying FIVE marks each and students have to attempt ALL questions.
- 3. SECTION-C contains THREE questions carrying TEN marks each and students have to attempt any TWO questions.

SECTION-A

- 1. What is meant by chemical potential?
- 2. Partial molar volume can be positive or negative-Explain.
- 3. Define activity coefficient of an electrolyte with suitable mathematical expression.
- 4. How the ionic strength is related with activity coefficient?
- 5. Give the number of phases, components and degree of freedom for the following:

Ice, water and vapour in equilibrium

- 6. What is an invariant system? Give an example.
- 7. What is meant by term eutectic? State the condition in which two substances can form a simple eutectic.
- 8. Write down Ilkovic equation with meaning of different parameters.
- 9. What is an azeotrope mixture? Give example.
- 10. What is meant by half wave potential in polarography?

1 M-51142 (S39)-2532

SECTION-B

- 11. Derive Gibbs-Duhem equation.
- 12. Derive an expression for the entropy of a mixture of ideal gases.
- 13. Calculate the total ionic strength of solutions that contains 0.30 M CaCl₂, 0.30 M Na₃PO₄, 0.10 M of Na₂SO₄ and 0.2 M NaCl.
- 14. Sketch well labelled phase diagram of water system.
- 15. Write a short note on primary salt effect.
- 16. Show that in the case of a mixture of ideal gases, the chemical potential of any constituent is given by the expression:

 $\mu_i = \mu_i(p)^0 + RT$ In p_i , where the symbols have their usual meaning.

SECTION-C

- 17. Draw and discuss the phase diagram for a three component system consisting of two solids A and B and water.
- 18. Show that for an ideal solution containing two components A and B, the Gibbs free energy of mixing is minimum when the mole fractions of the two components are the same, *i.e.*, equal to half each.
- 19. Describe the Lindemann theory of unimolecular reactions. Write a short note on flash photolysis for studying kinetics of fast reactions.

2 M-51142 (S39)-2532