

Roll No. 

--	--	--	--	--	--	--	--	--	--	--	--

Total No. of Pages : 02

Total No. of Questions : 07

**M.Sc Mathematics (2017 Batch) (Sem.-1)**

**MATHEMATICAL METHODS**

**Subject Code : MSM-105**

**Paper ID : [74724]**

**Time : 3 Hrs.**

**Max. Marks : 80**

**INSTRUCTION TO CANDIDATES :**

1. **SECTION-A is COMPULSORY** consisting of **EIGHT** questions carrying **TWO** marks each.
2. **SECTION - B & C.** have **THREE** questions in each section carrying **SIXTEEN** marks each.
3. **Select atleast TWO questions from SECTION - B & C EACH.**

**SECTION-A**

**1. Answer briefly :**

- a. Find the inverse laplace transform of  $\frac{s}{s^4 + s^2 + 1}$ .
- b. State the convolution theorem.
- c. Establish a relationship between fourier and laplace transforms.
- d. Enlist some applications of transforms to boundary value problems.
- e. Find the Z transform and radius of convergence of  $f(n) = 2^n, n < 0$
- f. Show that the geodesics on a plane are straight curves.
- g. Prove that the sphere is the solid figure of revolution in which given surface area has maximum volume.
- h. Define Kernal of the integral equation.

### SECTION-B

2.
  - a. Find the Laplace transform of  $\sin 2t \sin 3t$ .
  - b. Find the inverse transform of  $\frac{s^2 - 3s + 4}{s^3}$ .
3.
  - a. Define convolution of two functions  $f(x)$  and  $g(x)$  over the interval  $(-\infty, \infty)$  and Convolution theorem for Fourier transforms.
  - b. Find the Fourier cosine transform of  $e^{-x^2}$
4. Find the Z transforms of the following :
  - a.  $(n + 1)^2$
  - b.  $\sin (3x + 5)$
  - c.  $\cosh n\theta$
  - d.  $ne^{an}$

### SECTION-C

5. Solve the boundary value problem  $y'' - y' + x = 0$  ( $0 \leq x \leq 1$ ),  $y(0) = y(1) = 0$  by Rayleigh Ritz Method.
6. Use Galerkin's method to solve the boundary value problem which claims that the curve which extremizes the functional  $I$  such that;  
$$I = \int_0^{\pi/4} (y''^2 - y^2 + x^2) dx$$
 under the condition  $y(0) = 0$ ,  $y'(0) = 1$ ,  $y(\pi/4) = y'(\pi/4) = 1/\sqrt{2}$  is  $y = \sin x$ . Compare the approximate solutions with exact solutions.
7. Transform the differential equation  $y'' + y = x$ ,  $y(0) = 1$ ,  $y'(1) = 0$  to a fredholm integral equation, finding the corresponding Green's function.